


Safety Signal Observations

adverse events associated with different vaccines

By Craig Paardekooper



These observations are made using “Safety Signal” – an online search engine that ranks different vaccines by adverse effects.

Observations ([Check these yourself](#))

1. VACCINES & AUTISM

The vaccines with safety signals for autism are MEA, MMR, PNC, DTAP, HIBV and HEP – remarkably **5 of these are on the childhood vaccination schedule**. The two highest are MEA (the measles vaccine), and MMR (the measles, mumps and rubella vaccine). These vaccines also have safety signals for many other symptoms of autism.


There are a constellation of symptoms associated with autism, and each of these 5 vaccines have safety signals for most of them – **autism, autism spectrum disorder, antisocial behavior, neurodevelopmental disorder, Subacute sclerosing panencephalitis, asperger, social avoidant behaviour, mental retardation, decreased eye contact, abnormal behaviour, obsessive compulsive disorder, hostility, personality disorder, psychomotor hyperactivity, developmental delay, faecal incontinence, speech disorder developmental, screaming, attention deficit disorder, developmental regression, stupor, aggression, febrile convulsion, childhood disintegrative disorder, learning disability, regressive behaviour, disturbance in social behaviour, convulsion, staring, hearing impaired, intentional self-injury.**

CHILDHOOD IMMUNIZATIONS – COMBO 10	
Product lines	BCN Commercial
Source	HEDIS
Description	<p>The percentage of children 2 years of age who meet the combination 10 criteria on or before their second birthday:</p> <ul style="list-style-type: none">• (4) DTaP* vaccinations• (3) IPV* vaccinations• (1) MMR vaccination• (1) VZV vaccination• (3) HiB* vaccinations• (3) Hepatitis B vaccinations• (4) PCV* vaccinations• (1) HepA vaccination• (2 or 3) RV* vaccinations• (2) Influenza** vaccinations <p>*Vaccinations administered prior to 42 days after birth are not counted as a numerator hit. **Vaccinations administered prior to 180 days after birth are not counted as a numerator hit.</p>
Continuous enrollment	Must be continuously enrolled 12 months prior to child's second birthday

Note that these 5 vaccines are unique in being the only vaccines that generate safety signals for **SIDS (sudden infant death syndrome)** ! [Check it yourself](#) !

All 5 are on the childhood schedule. The effect of all 5 vaccines on an infant would be to compound the probability of brain damage. Even when clinical levels of brain damage are absent, there may still be sub-clinical levels, causing permanent cognitive deficit.

Does this explain the steady rise in autism since 2000 A.D.



2. VACCINES & VASCULAR DISORDER

Vascular disorders include **thrombosis, infarction, embolism, aneurysm, ischaemia, bleeding, haemorrhage, haematoma, menorr*, arteriosclerosis, phlebitis** . Only COVID vaccines have safety signals for these symptoms.

The essential nature and predominant characteristic of COVID vaccines is their strong, and unique association with vascular disorder. Please see [here](#).

3. VACCINES & CARDIAC DISORDERS

Symptoms include **myocarditis, pericarditis, tachycardia, arrhythmia, bradycardia, chest pain, palpitations, flutter, atrial *, aortic *, cardiac arrest/failure, sudden death**.

The vaccines with these safety signals include COVID19 and SMALLPOX

Cardiac failure/death/arrest is only associated with COVID vaccines. Hence "sudden death". Its almost as if COVID stands for **CO**ronary **V**ascular **I**schaemic **D**isease. Please see [here](#).

4. VACCINES & LIVER INJURY

Symptoms include – **liver disorder, liver tenderness, acute hepatitis B, autoimmune hepatitis, hepatic cirrhosis, hepatic enzyme abnormal, hepatic enzyme increased, hepatic failure, hepatic function abnormal, hepatic necrosis, hepatic steatosis, hepatitis, hepatitis A, hepatitis B, hepatocellular damage, hepatomegaly, hepatosplenomegaly, viral hepatitis carrier**.

If you were asked “which vaccine is associated with these disorders?” you would probably not suggest HEP, since HEP vaccines are supposed to protect us from liver disorder. However HEP vaccines are EXCLUSIVE in having safety signals for ALL of these disorders. Liver / hepa* disorders/ injury are unique to hepatitis vaccines (HEP) ! [Check it yourself](#) !

5. VACCINES & BRAIN INFLAMMATION

Symptoms include **encephalitis, central nervous system inflammation, acute disseminated encephomyelitis, autoimmune encephalopathy, encephalopathy, encephalogram abnormal, autoimmune encephalitis**. The vaccines with safety signals for encephalitis are MEA, MMR, YF, HEP, PNC, DTP, HPV, FLUX. Central Nervous System inflammation is associated with HPV, HEP and FLUX. Encephalitis is most strongly associated with MMR and MEA, though also associated with HPV, HEP and FLUX.

Cerebral Autoimmune diseases are most strongly associated with HPV

6. VACCINES & BODY-WIDE INFLAMMATION

Symptoms include **multisystem inflammation syndrome in adults, multisystem inflammatory syndrome in children**.

Only COVID vaccines have safety signals for MIS. This is because LNPs carry the vaccine mRNA to all parts of the body, and spike protein is generated in all parts. Generation of spike on cell surfaces causes autoimmune attack on those cells and resulting systemic inflammation. [Check it yourself](#) !

Multisystem inflammation syndrome kills.

7. VACCINES & CANCER

Symptoms include **cancer, neoplasm, neoplasia, metast***, tumour, lymphoma, leukaemia, carcinoma

Cancer : Covid, HPV

Neoplasm : Covid, OPV,

Metast* : Covid.

Tumor : Covid

Lymphoma : HEP

Leukaemia : HPV4, MMR

Carcinoma : HPV4, Covid,

HPV4 and Covid vaccines are the predominant vaccines with safety signals for cancer. [Check it yourself](#) !

8. TREMOR

Symptoms include tremor, muscle twitching, convulsion, epilepy, spasm

Only DTP and DTPIPV have significant safety signals for tremor

DTP has the highest safety signal for grand mal convulsion

DTP is also associated with encephalitis, so perhaps encephalitis causes the tremor.

Only HPV has safety signals for tonic convulsion

Only HPV4 has a safety signal for juvenile myoclonic epilepsy

HPV, MMR and DTP have safety signals for epilepsy

MENB, HPV and MMR have safety signals for petit mal epilepsy

[Check it yourself](#) !

9. INFECTION

Only VARZOS and VARCEL have safety signals for Herpes

Only COVID vaccines have safety signals for Covid

Only COVID vaccines have a safety signal for Covid pneumonia

Only HEP vaccines have safety signals for Hepatitis

Only MMR has safety signals for Mumps

Only MMR has safety signals for Rubella

Only RV1,RV5, RVX have safety signals for Rotavirus infection

Only MEA and MMR have safety signals for measles

Only HPV has safety signals for human papilloma virus

TDAPIPV, PER have the highest safety signals for Pertussis (Whooping cough)

PNC, PPV have the highest safety signal for pneumonia

FLU vaccines have the highest safety signals for influenza

Only VARCEL, VARZOS and MMRV have safety signals for varicella virus (Chickenpox)

Only TDAP, COVID vaccines have a safety signal for HIV antibody

Only HEP has a safety signal for HIV test positive

MEN, MNQ vaccines have a high safety signal for meningitis

HIBV, PNC, MMR, HEPAB (childhood vaccines) have safety signals for meningitis

[Check it yourself](#) !

This is an interesting phenomenon. The pattern seems to be “a vaccine induces the illness that it is meant to protect from, far more than other vaccines”.

This data suggests that vaccinated are more prone to illness than unvaccinated, and this requires investigation. If this effect is not attributable to a reporting bias, then it could be due to direct injection of a live pathogen or to immunosuppression or both.

10. ALLERGIES

Allergies - eczema, food allergies, asthma, anaphylactic reaction, anaphylactic shock, anaphylactoid reaction

MMR, HPV4, RV1, RV5 have safety signals for food/milk allergies

COVID, DTAPIPVHIB, SMALL, MMR have safety signals for eczema

FLU, HEP, MMR have safety signals for asthma

HEP, HPV2, MMR, FLU, TYP, TD have safety signals for anaphylactoid reaction

DF, FLUX(H1N1), TTOX have safety signals for anaphylactic reaction

HEP, HPV2, RAB have safety signals for anaphylactic shock

11. KIDNEYS

Symptoms include acute kidney injury, chronic kidney disease, biopsy kidney abnormal, end stage renal disease, renal colic, renal failure, renal failure acute

Acute Kidney Injury : COVID, YF

Chronic kidney disease : COVID

Biopsy Kidney abnormal : FLUX, HPV4

End stage renal disease : COVID

Renal colic : COVID

Renal Failure : YF

Renal failure acute : FLUX, FLU3, HEP, PPV, HPV4

Renal infarct : COVID

Renal Pain : COVID

COVID and Yellow Fever vaccines have safety signals for kidney injury. This is also evident from an analysis of adverse reactions following COVID vaccination see [symptoms-delayed-immediate.pdf \(howbad.info\)](#). Note the ranking of acute kidney injury.

Vaccines are exempt from liability. [Why vaccine manufacturers are exempt from liability – Sonoran News](#) Maybe this needs to change, but until then more caution should be exercised. The above risks are never or very reluctantly acknowledged by pharmaceutical manufacturers or by the medical profession they sponsor.